Astro 404 Lecture 12 Sept. 23, 2019

Announcements:

• Problem Set 4 due next Friday

wordy but less mathematical than PS3 instructor office hours: Wed 11am-noon or by appt

• PS grades not posted yet, but will start to appear very soon

Last time:

• Ideal gas theory

Q: microscopic origin of pressure? temperature?

- Virial theorem
- ← Q: what's that? result for ideal gas?
 - *Q*: when does it apply?

Ideal Gas Recap

microscopic theory gives ideal gas pressure

$$P = N \frac{m_{\rm g} \langle v^2 \rangle / 3}{V} = n \ kT \tag{1}$$

- $P \propto N$: more particles \rightarrow collisions more frequent
- $P \propto 1/V$: more $V \rightarrow$ collisions less frequent
- $kT = m_g \langle v^2 \rangle / 3$: faster particles \rightarrow hotter gas

Virial Theorem Recap

for a *monatomic ideal gas* in *hydrostatic equilibrium* the Virial theorem relates global stellar energy reservoirs

$$U = -\frac{1}{2}\Omega\tag{2}$$

total internal energy is minus half total grav potential energy

a kind of stellar equipartition

like ideal gas kinetic equipartition: $m_g \langle v_x \rangle^2/2 = m_g \langle v_y \rangle^2/2 = m_g \langle v_z \rangle^2/2 = kT$

the total energy in a star under hydrostatic equilibrium is

$$E_{\text{tot}} = U + \Omega = \frac{1}{2}\Omega = -\frac{1}{2}\int \frac{Gm \ dm}{r}$$
(3)

ω

Q: order of magnitude for each?

Virial Theorem: Order of Magnitude

to order of magnitude:

- internal/thermal energy: $U \sim M \langle kT \rangle / m_g = N \langle kT \rangle$
- gravitational potential energy: $\Omega \sim -GM^2/R$

Virial theorem links these, and implies

$$kT \sim \frac{GMm_{\rm g}}{R} \tag{4}$$

note: $kT \sim m_{\rm g} \langle v^2 \rangle$, and $GM/R \sim \Phi$ so Virial also means that

$$\langle v^2 \rangle \sim \Phi$$
 (5)

▶ particle thermal speeds set by depth of gravitational potential

Virial Theorem and Stellar Evolution

note that Virial theorem relates

- global stellar energy reservoirs: gravitational and internal/thermal
- and hence total energy localized to the star
- while in state of *hydrostatic equilibrium*

but recall that stars lose energy – they are luminous!

Q: how can a star do this and maintain equilibrium state?

Maintaining Equilibrium: Burning Phases

Virial theorem: $\langle kT \rangle \sim GMm_{\rm g}/R$, and $E_{\rm tot} \sim -N \langle kT \rangle \sim -GM^2/R$

to maintain $E_{tot} \rightarrow$ must maintain R and T

- so the star must *tap some non-gravitational "fuel source"* that supplies energy to maintain *T* and power luminosity
- a given equilibrium state (R, T, L) lasts as long as fuel permits

Thus we anticipate **burning phases set by fuel supplies**

Q: when will a star change its equilibrium state?

σ

Seeking Equilibrium: Contraction Phases

if the "fuel source" is gravitational energy itself

radiated energy *decreases* E_{tot} , i.e., makes it *more negative*

- \bullet Virial theorem demands that R decreases as well
- the star contracts!
- and in response *heats up!*

this continues unless/until a new fuel source ignites

Thus we expect **contraction phases** between burning phases

Perturbing a Star

consider a star maintaining hydrostatic equilibrium by burning a non-gravitational fuel supply to generate energy

now imagine a *perturbation that generates energy faster* due to upward density or temperature fluctuation at the star's center ("core")

if the star is composed of an *ideal gas*:

- Q: how does the star's center respond?
- *Q*: what is the effect on the fuel burning rate?
- *Q*: what is the net response to the perturbation?

00

Q: what if a perturbation had lowered the core burning rate?

The Virial Theorem and Stellar Stability

an increased core energy production rate:

- ▷ leads to a higher core temperature
- ▷ for ideal gas: increases pressure
- ▷ pressure gradient drives *core expansion*
- b density drops, pressure and temperature drop
- ▷ this *lowers burning rate* until
- ▷ the star recovers its initial state!

can convince yourself:

same conclusion for downward perturbation

Lesson: perturbed ideal gas stars burning non-gravitational fuel

are driven back to initial state
the equilibrium is stable!

The Virial Theorem and Stellar Contraction argument courtesy of Alessandro Chieffi

consider a star not generating energy via "fuel burning" and thus contracting

if contraction isn't too fast

star passes through a series of states near hydrostatic equilibrium in which total energy is

$$E_{\text{tot}} = -U_{\text{int}} \sim -N \ \langle kT \rangle \tag{6}$$

transition between states requires change of internal energy this takes time to occur

⁶ "protects" the star against sudden violent changes

iClicker Poll: Ultra-Relativistic Stars

so far, implicitly assumed stars are *non-relativistic* ideal gasses

Consider a star composed of a gas of *relativistic particles* that is: speeds $v \approx c$ or even v = c

Vote your conscience!

Will the star be more or less stable than if non-relativistic?

A more stable (faster = more internal energy = more tightly bound)

less stable (faster = more pressure = less tightly bound)

11

С

no change in stability (effects cancel)

Ideal Gas: Ultra-Relativistic Case

thus far implicitly assumed *gas speeds are non-relativistic*

 \bullet speeds $v \ll c$ and $p \ll m_{\rm g}c$

12

• so $p^2/m_{\rm g} \ll m_{\rm g}c^2$ and thus $kT \ll m_{\rm g}c^2$

now consider opposite limit: ultra-relativistic particles

- $v \approx c$ or even v = c Q: examples?
- relativistic momentum $p \gg m_{\rm g}c$
- energy $E = \sqrt{(cp)^2 + (m_g c^2)^2} \approx cp$

revisit microscopic pressure derivation

$$P_{\text{rel}} = \frac{N \langle pv \rangle_{\text{rel}}}{3V} \approx \frac{1}{3} \frac{N \langle E \rangle}{V} = \frac{1}{3} \frac{U_{\text{rel}}}{V}$$
(7)

and thus for a relativistic gas PV = U/3, and energy density is

$$\varepsilon_{\rm rel} = 3P_{\rm rel}$$
 (8)

Virial Theorem for an Ultra-Relativistic Gas

basic Virial theorem argument still holds

$$\int P \ dV = -\frac{1}{3}\Omega_{\text{grav}} \tag{9}$$

but for relativistic gas: pressure and energy density related via $P_{\text{rel}} = \varepsilon_{\text{rel}}/3$ instead of non-relativistic $P_{\text{nr}} = (2/3)\varepsilon_{\text{nr}}$ so $\int P_{\text{rel}} dV = (1/3) \int \varepsilon_{\text{rel}} dV = U_{\text{rel}}/3$

thus for a relativistic gas, Virial theorem is

$$U_{\rm rel} = -\Omega_{\rm grav} \tag{10}$$

Q: so what is total energy? implications?

 $\tilde{\omega}$ Q: for a fixed mass distribution = fixed Ω , which gas is hotter?

Ultra-Relativistic Stars are Unstable

relativistic gas in equilibrium: Virial theorem

$$U_{\rm rel} = -\Omega_{\rm grav} \tag{11}$$

this gives total energy

$$E_{\text{tot}} = \text{internal} + \text{grav pot} = U_{\text{rel}} + \Omega = 0$$
 (12)

total energy is zero!

dramatic implications:

- $E_{tot} = 0$ means system is marginally bound
- transition between equilibrium states requires no change in internal energy
- star can evolve violently: **a relativistic star is unstable!**

Virial Theorem: Lessons

equilibrium links thermal and gravitational energy more compact \leftrightarrow hotter

- stellar interiors much hotter than $T_{\rm eff}$
- as stars lose energy they get hotter!

15

- (non-relativistic) ideal gas stars are self-regulating: stable
- (non-relativistic) ideal gas stars require time to evolve
- relativistic stars are barely bound, can evolve rapidly these stars are unstable!

Stars: Energy Generation

How Does the Sun Shine?

The Sun radiates: shines from thermal radiation

• recall: surface flux
$$F_{surf,\odot} = \sigma T_{surf,\odot}^4 = 60 \text{ MWatt/m}^2$$

- total power output = rate of energy emission = luminosity $L_{\odot} = 4\pi R_{1}^{2} _{AU}F_{\odot}(1 \text{ AU}) = 3.85 \times 10^{26} \text{ Watts} \qquad (13)$ $\rightarrow \text{ the Sun is a } 4 \times 10^{26} \text{-Watt lightbulb}$
- But also: the Sun has *constant* temperature, luminosity (over human timescales \gtrsim centuries)
- $\stackrel{\vdash}{\neg}$ Q: how is the Sun unlike a cup of coffee?

The Sun is Not a Cup of Coffee

Coffee Thermodynamics

Demo: cup of coffee: cools thermodynamic lesson:

- left alone, hot coffee cools (surprise!) \rightarrow energy radiated, not replaced
- to keep your double-shot soy latte from cooling need Mr. CoffeeTM machine–energy (heat) source

Contrast with the Sun

- surface T_{\odot} constant over human lifetimes but energy *is* radiated, at enormous rate
- ergo: something must replace the lost energy
- ▷ What is solar heat source (fuel supply)?
 - \rightarrow a mystery in Astronomy until the 20th century
- $\overline{5}$ Q: all possible energy/heat sources which Sun taps? Q: how to test/compare which are important?

Energy Conservation and the Sun

recall: power is energy flow rate L = dE/dt

assume:

- Sun always emits energy at today's rate (L constant)
- radiation lasts for time τ_{\odot} = "lifetime" of Sun Q: what is a minimum value for τ_{\odot} ?

energy output over Sun's lifetime:

 $E_{\text{lost}} = L\tau$

Energy conservation:

solar energy supply = lifelong energy output

19

Solar Batteries: Required Lifetime

we found from radioactive dating of meteorites: the solar system is very old: age $t_{SS} = 4.55 \times 10^9 \ yr$ Sun's present age essentially the same: $t_{\odot,now} = t_{SS} = 4.55$ billion years

total energy output over this time is huuuge! \rightarrow required huge energy reservoir

in other words: important solar energy source(s) \equiv long-lived: $\tau_{\text{source}} = E_{\text{res}}/L_{\odot} = \tau_{\odot} > t_{\odot,\text{now}} \approx 5$ billion yr

Q: possible sources-not just right answer, but any energy reservoirs?

iClicker Poll: Rank the Energy Sources

Vote your conscience!

Of the proposed solar energy reservoirs

Which one is the largest, i.e., can power the Sun longest?

Which one is the smallest?