

Astro 404  
Lecture 31  
Nov. 8, 2019

Announcements:

- **Problem Set 9 due today**

typo corrected in L24 notes:  $n_Q = (2\pi mkT/h^2)^{3/2}$

- **Problem Set 10 due next Friday Oct 15**

Last time: down with solar-mass stars! onward to massive stars!

*Q: what for us are high-mass stars?*

## Massive Star Demographics

in our context, massive:  $M \gtrsim 8 - 10M_{\odot}$   
that is: destined to become core-collapse supernovae

PS10: study **initial mass function**

distribution of star birth masses

- massive stars are  $\sim 0.5\%$  by *number* of all stars born
- but comprise  $\sim 10\%$  of *mass* going into stars

*Q: how can these both be true?*

lesson: massive stars are rare but spectacular celebrities of the cosmos

## Massive Stars: Radiation Pressure

radiation force on electron with cross section  $\sigma_e$  (PS1):

$$F_{\text{rad}} = P_{\text{rad}}\sigma_e = \frac{L\sigma}{4\pi r^2 c} \quad (1)$$

inverse square law! same as gravity but repulsive!

*radiation force balances gravity* on  $e + p$  pair when

$$L = L_{\text{Edd}} = \frac{4\pi GMm_p c}{\sigma} \quad (2)$$

*Eddington luminosity*

Q: what if  $L > L_{\text{Edd}}$ ?

# Massive Stars and the Eddington Luminosity

*Eddington luminosity*:  $F_{\text{rad}} = F_{\text{grav}}$  when

$$L = L_{\text{Edd}} = \frac{4\pi GMm_p c}{\sigma} \quad (3)$$

if  $L > L_{\text{Edd}}$ : radiation pressure stronger than gravity!  
star pushes its own atmosphere away

→ Eddington gives *maximum stable luminosity*

PS7: massive stars have  $L$  very near  $L_{\text{Edd}}$ !

- near the edge of stability!
- drive strong winds even during main sequence
- mass loss important (and uncertain) over entire star life

5

Q: *consequences of strong mass loss?*

# The Highest(?) Masses: Wolf-Rayet Stars

for the very *highest masses*:  $M \gtrsim 30M_{\odot}$ ?

and with solar composition

- ★ *mass loss very strong* even in main sequence
- ★ *reduces star mass* → converge to  $30M_{\odot}$ ?
- ★ hydrogen envelope can be completely removed and *helium core exposed* (and sometimes deeper)
- ★ wind material shows nucleosynthesis products e.g., CNO cycle abundance pattern: nitrogen rich

observed at *Wolf-Rayet* stars

www: Wolf-Rayet wind

◦ eta Carinae: initially  $120M_{\odot}$ ? now  $\sim 100M_{\odot}$

www: eta Carinae

## iClicker Poll: Massive Stars on the HR Diagram

evolution drives  $L \rightarrow L_{\text{Edd}} \propto M$

Implications for a given mass on HR diagram?

- A** HR evolution nearly horizontal
- B** HR evolution nearly vertical
- C** HR evolution keeps  $L/T_{\text{eff}}$  fixed

## Massive Stars on the HR Diagram

evolution drives  $L \rightarrow L_{\text{Edd}} \propto M$

also recall: main sequence is sequence in mass  
so on main sequence, for all stars:  $L$  grows with mass

and for massive stars:

$L \rightarrow L_{\text{Edd}}$  fixed by mass (roughly) on MS and beyond

so *post-main-sequence evolution changes  $T_{\text{eff}}$  but not  $L$*   
→ motion on HR diagram is horizontal

## Massive Stars: Burning Phases

Main sequence: hydrogen burning

- convective core  $\rightarrow$  fuel circulation
- $T_c \gtrsim 2\times$  hotter than Sun
- burn  $p \rightarrow {}^4\text{He}$  via CNO cycle  
avoid Weak  $pp \rightarrow de\nu$ : goes much faster

*when core hydrogen exhausted:*



core contracts, smoothly begins burning helium

non-degenerate, no helium flash

with hydrogen burning in shell


star becomes a **supergiant**

www: Betelgeuse imaged







# Massive Stars on the HR Diagram: Supergiants



When core He exhausted, begin cycles:

- **contract**
  - ignite new **shell burning**
  - **ignite ash** → fuel in core
  - **burn core to exhaustion**
- repeat...

develop “onion skin” structure: **www: pre-SN**  
 favors “ $\alpha$ -elements” : tightly bound



## Neutrino Cooling

At  $T \gtrsim 5 \times 10^8$  K (C burn):

neutrinos produced via  $e^+e^- \rightarrow \nu\bar{\nu}$

much slower than  $e^+e^- \rightarrow \gamma\gamma$  yet still crucial

Q: *why?*

neutrino production rate per volume:

$$q_\nu = \langle \sigma v n_e^2 \rangle \sim T^2 \times (T^3)^2 \sim T^8 \quad (4)$$

$\nu$  escape  $\rightarrow$  dominate  $E$  loss: **neutrino cooling**

neutrino  $E$  loss rate per vol:  $\varepsilon_\nu = E_\nu q \sim T^9$

equilibrium:  $\varepsilon_{\text{emit},\nu} = \varepsilon_{\text{released,nuc}}$

$\rightarrow L_\nu \sim (1 - 10^{-6})L_\gamma$  for C thru Si burning: **neutrino star!**

## iClicker Poll: Effect of Neutrino Losses

when neutrino emission dominates total luminosity:

What is effect on burning phases?

- A** neutrino star burning phases last a *longer* time than if no neutrinos emitted
- B** neutrino star burning phases last a *shorter* time than if no neutrinos emitted
- C** neutrino star burning phases last the *same* time than if no neutrinos emitted

## Si Burning

neutrino emission removes energy from core  
“steals” nuclear energy now unavailable to heat star  
shortens burning phases—final stages: months, days

$T \sim 4 \times 10^9$  K  $\rightarrow$  photon energy density  $\epsilon_\gamma \sim T^4$  large  
photodisintegration  $^{28}\text{Si} + \gamma \rightarrow p, n, \alpha$

1.  $\gamma$ s take  $p, n, \alpha$  from weakly bound nuclei
2. these recombine with all nuclei
3. flow  $\rightarrow$  more tightly bound

Net effect: redistribute to most tightly bound nuclei

## Binding Energy Patterns

recall: binding energy  $B_i$  is  
energy required to tear nucleus to protons and neutrons

note that larger nuclei have large  $B_i$ ,  
but shared among more nucleons


consider: **binding energy per nucleon**  $B/A$

*Q: what does this represent physically?*

# Nuclear Stability: Binding Energy

For stable nuclei:

- sharp rise in  $B_i/A_i$  at low  $A$
- local max at  ${}^4\text{He}$
- *no stable nuclei at  $A = 5, 8$*
- lowest  $B/A$  for D, LiBeB
- *max  $B/A$  for middle masses:*
- **peak at  ${}^{56}\text{Fe}$**



## Nuclear Equilibrium

nuclear reactions drive core to **equilibrium**  
dominated by most stable nuclei possible  
→ most tightly bound

max abundance → largest nuclear binding: “iron peak”

*core dominated by iron and nickel*

And now the end is imminent. Q: *why?*



## Iron Core Evolution

*can't burn Fe → degenerate core*

support:  $e$  degeneracy pressure—core is iron white dwarf!

first time a massive star core is degenerate

stable briefly, but...

do burn Si in overlying shell


→ increase Fe core mass

when  $M_{\text{core}} > M_{\text{Chandra}}$  → core unstable

begins to collapse

## Core Collapse

upon collapse: *iron core disintegrated by photons*



huge density: electrons have high Fermi energy  
→ favorable to get rid of them!

**electrons capture** onto protons  $e^- + p \rightarrow n + \nu_e$

and onto nuclei  $e^- + Z_A \rightarrow Z - 1_A + \nu_e$

“neutronization” or “deleptonization”

removes  $e$  and so reduces degeneracy pressure!

- accelerates collapse (positive feedback)
- also: releases  $\nu_e$

## Collapse Dynamics

*Freefall timescale* for material with density  $\rho$  (PS4):

$$\tau_{\text{ff}} \sim \frac{1}{\sqrt{G\rho}} \sim 446 \text{ s} \sqrt{\frac{1 \text{ g/cm}^3}{\rho_{\text{cgs}}}} \lesssim 1 \text{ sec}$$

but pre-supernova star very non-uniform density

*Q: what does this mean for collapse?*

inner core: homologous collapse  $v \propto r$

outer core: quickly becomes supersonic  $v > c_s$

outer envelope: unaware of collapse

*Q: what (if anything) stops collapse?*

## Bounce and Explosion

core collapses until  $\rho_{\text{core}} > \rho_{\text{nuc}} \sim 3 \times 10^{14} \text{ g/cm}^3$

repulsive short-range nuclear force dominates: *“incompressible”*

details depend on equation of state of nuclear matter

1. *core bounce* → proto neutron star born
2. *shock wave* launched
3. a miracle occurs
4. outer layers *accelerated*

*Demo: AstroBlaster™*

5. successful *explosion* observed  
→  $v_{\text{ej}} \sim 15,000 \text{ km/s} \sim c/20!$

**Why step 3?** What's the miracle?

“prompt shock” fails:

do launch shock, but

- overlying layers infalling

→ ram pressure  $P = \rho v_{\text{in}}^2$

- dissociate Fe → lose energy

shock motion stalls → “accretion shock”

“prompt explosion” mechanism fails

*Q: what needed to revive explosion?*

## Delayed Explosion Mechanisms

“delayed explosion” to revive:

neutrinos, 3-D hydro/instability, rotation effects?

some models not work, but controversial

**Energetics:**

$$E_{\text{ejecta}} \sim M_{\text{ej}} v^2 \sim (10 M_{\odot}) (c/20)^2 \sim 10^{51} \text{ erg} \equiv 1 \text{ foe}$$

but must release gravitational binding energy

$$\begin{aligned} \Delta E &\sim -GM_{\star}^2/R_{\star} - (-GM_{\text{NS}}^2/R_{\text{NS}}) \\ &\simeq GM_{\text{NS}}^2/R_{\text{NS}} \sim 3 \times 10^{53} \text{ erg} = 300 \text{ foe} \end{aligned}$$

*Q: Where does the rest go?*

⇒ SN calculations must be good to  $\sim 1\%$

to see the minor optical fireworks