

Astro 404  
Lecture 32  
Nov. 11, 2019

Announcements:

- **Problem Set 10 due next Friday Oct 15**
- Office Hours: Instructor – after class or by appointment
- TA: Thursday noon-1pm or by appointment
- Exam: grading elves hard at work

Last Time: core-collapse supernovae-prelude to explosions

*Q: core-collapse progenitors: masses? lifetimes?*

*Q: main seq location HR diagram? evolution?*

*Q: nuclear burning phases? nucleosynthesis products?*

↳ *Q: neutrino production—during which phases? Origin?*

*Q: evolution after main sequence? core structure?*

massive stars:  $8 - 10M_{\odot}$   
“celebrities of the cosmos”

- live fast: high  $T_C, \rho_C$   
→ rapid nuclear burning
- die young:  
lifetimes  $\sim$  few Myr
- we’ll see: leave beautiful corpse


## Massive Star Binarity

recall that most stars overall are in binaries

★ nearly 100% of massive stars are in binaries

★ often the binary companion is another massive star!


this fact will be important



after main sequence: repeated cycles of

- core contraction and ignition
- ash of last burning phase becomes fuel for next
- shell burning “remembers” earlier phases

develop “onion skin” structure:  $\omega$  pre-SN  
favors “ $\alpha$ -elements” : tightly bound



## Collapse Dynamics

*Freefall timescale* for material with density  $\rho$  (PS4):

$$\tau_{\text{ff}} \sim \frac{1}{\sqrt{G\rho}} \sim 446 \text{ s} \sqrt{\frac{1 \text{ g/cm}^3}{\rho_{\text{cgs}}}} \lesssim 1 \text{ sec}$$

but pre-supernova star very non-uniform density

*Q: what does this mean for collapse?*

inner core: homologous collapse  $v \propto r$

outer core: quickly becomes supersonic  $v > c_s$

outer envelope: unaware of collapse

‡ *Q: what (if anything) stops collapse?*

## Bounce and Explosion

core collapses until  $\rho_{\text{core}} > \rho_{\text{nuc}} \sim 3 \times 10^{14} \text{ g/cm}^3$

repulsive short-range nuclear force dominates: *“incompressible”*

details depend on equation of state of nuclear matter

1. *core bounce* → proto neutron star born
2. *shock wave* launched
3. a miracle occurs
4. outer layers *accelerated*

*Demo: AstroBlaster™*

5. successful *explosion* observed  
→  $v_{\text{ej}} \sim 15,000 \text{ km/s} \sim c/20!$

**Why step 3?** What's the miracle?

“prompt shock” fails:

do launch shock, but

- overlying layers infalling

→ ram pressure  $P = \rho v_{\text{in}}^2$

- dissociate Fe → lose energy

shock motion stalls → “accretion shock”

“prompt explosion” mechanism fails

*Q: what needed to revive explosion?*

## iClicker Poll: Supernova Neutrinos

We saw that the Sun is a confirmed source of neutrinos in fact: a few percent of the Sun's luminosity (energy release) is in neutrinos rather than light

Now consider a massive star, exploding as a supernova and vote your conscience:

Which best describes a supernova's energy release?

**A** < 1% of energy released in neutrinos, > 99% in photons

**B**  $\approx$  50% of energy released in neutrinos,  $\approx$  50% in photons

2 **C** > 99% of energy released in neutrinos, < 1% in photons

## Delayed Explosion Mechanisms

“delayed explosion” to revive:

neutrinos, 3-D hydro/instability, rotation effects?

some models do work, but controversial

**Energetics:**

$$E_{\text{ejecta}} \sim M_{\text{ej}} v^2 \sim (10 M_{\odot}) (c/20)^2 \sim 10^{51} \text{ erg} \equiv 1 \text{ foe}$$

but must release gravitational binding energy

$$\begin{aligned} \Delta E &\sim -GM_{\star}^2/R_{\star} - (-GM_{\text{NS}}^2/R_{\text{NS}}) \\ &\simeq GM_{\text{NS}}^2/R_{\text{NS}} \sim 3 \times 10^{53} \text{ erg} = 300 \text{ foe} \end{aligned}$$

*Q: Where does the rest go?*

⇒ SN calculations must be good to  $\sim 1\%$

∞

to see the minor optical fireworks



# Supernova Neutrinos

two phases of neutrino emission during collapse and explosion:

1. **neutronization**
2. **thermal emission**

when electrons removed to make neutrons  
neutronization neutrinos produced before collapse  
emitted over  $< 1$  sec, leave freely

during collapse: huge temperature  $kT > m_e c^2$   
thermal bath makes  $e^+e^-$  pairs  
sometimes make **thermal neutrinos**  $e^+e^- \rightarrow \nu\bar{\nu}$

## Thermal Supernova Neutrinos

by far, thermal neutrinos have a larger luminosity and larger energies than neutronization neutrinos  
→ these are the bulk of the supernovae neutrino emission

thermal  $\nu$ s initially leave freely  
but when proto-neutron-star formed  
mean free path  $l_\nu = 1/(n_{\text{nuc}}\sigma_\nu)$   
becomes small:  $l_\nu \lesssim R_{\text{NS}}$

*Q: what happens to these thermal neutrinos?*

*Q: will they ever escape? if so, how?*

*Q: neutrino telescope time signature?*

## Supernova Neutrinos: Theory

when dense core has  $\ell_\nu \lesssim R_{\text{NS}}$ : neutrinos trapped  
proto-neutron star develops “neutrinosphere”  
size set by radius where  $\sim 1$  scattering to go:  $r \sim \ell_\nu(r)$

inside  $r_\nu$ : weak equilibrium  $\rightarrow$  “neutrino star”

- both neutrinos and anti-neutrinos created  
for experts: all species  $\nu_e, \nu_\mu, \nu_\tau \approx$  equally populated

neutrinos still leave, but must diffuse

emit neutrinos & energy (cool) over diffusion time

PS10:  $\tau_{\text{diff}} \sim \text{few sec}$

- ≡ Q: *how to test this? how to find supernovae? where to look?*  
Q: *how to identify progenitor (pre-explosion star)?*

# Supernovae Observed: Historical Supernovae

*supernovae are rare:*

- true rate: about  $\sim 3/\text{century}$  in our Galaxy
- observed (naked-eye) rate:  $\sim 0.5/\text{century}$   
our Galaxy dims and obscures most supernovae!

## Supernovae Discovery Strategy I:

*look at written records in historical archives*

try to match with known explosion remnants on sky

**pro:** get firsthand account!

**con:** ancient records often ambiguous

and no hope of learning about pre-supernova (progenitor) star

# Supernova 1054

- July 4(!) 1054: event seen in Taurus
- no record in Europe, even though should have been visible
- “guest star” noted in Chinese astronomical records
- also possible hint in Anasazi (Pueblos) rock paintings  
www: Anasazi drawing, Y1K
- possible indications in artifacts from India
- Present-day: Crab Nebula (Messier 1)  
www: present-day view: Y2K  
one of the closest and best-studied supernova remnants!

## Supernova 1572

reported extensively by Tycho Brahe: “Nova Stella” – new star

www: sketch

On the 11th day of November in the evening after sunset ... I noticed that a new and unusual star, surpassing the other stars in brilliancy, was shining ... and since I had, from boyhood, known all the stars of the heavens perfectly, it was quite evident to me that there had never been any star in that place of the sky ...

I was so astonished of this sight ... A miracle indeed, one that has never been previously seen before our time, in any age since the beginning of the world.

– Tycho Brahe

*Q: What did Tycho get right? Where was he wrong?*

## Tycho's Supernova

- ★ Tycho recorded brightness peaked after days then visible for months
- ★ Searched for but did not find **parallax** showed event had to be at a great distance certainly beyond the Moon
- ★ dramatic challenge to Aristotelian/Ptolemaian worldview celestial realm supposed to be perfect and unchanging: “incorruptible” very different from “corruptible” terrestrial realm we live in Tycho showed the heavens are changeable

# Extragalactic Supernovae

## *Supernova Detection Strategy II*

since only a few per century per galaxy, *look at many galaxies!*

→ if monitor 100 Milky-Way-like galaxies,  
expect to see  $\sim$  *few* supernovae per year!

**pro:** much higher discovery rate

if know distance to galaxy, get distance to SN

can find events with little dust obscuration

can search for progenitor stars in archival images

**con:** don't know where or when a supernova will occur

must monitor many galaxies over a long time

farther away → less able to resolve details

this has been incredibly successful:

**most of our SN knowhow comes from extragalactic events**



# Observed Supernovae: Properties and Correlations

**spectra** of supernovae after explosions show two classes

**Type I: hydrogen** totally or nearly *absent*

in spectrum and thus ejecta

subclasses: Type Ia: silicon present, iron-peak elements

Types Ib and Ic: helium and oxygen present

**Type II: hydrogen present** in spectrum and ejecta

*Q: how could we understand this?*

**host galaxies** show correlation with type

*elliptical/early-type galaxies*: no/little ongoing star formation

- only have Type Ia explosions
- no progenitors identified

*spiral and irregular galaxies*: star formation ongoing

- supernovae found in star-forming regions
- Types Ib, Ic, and II all found
- progenitors have masses  $8 - 50M_{\odot}$
- Type Ib and Ic progenitors:  
evidence of winds, Wolf-Rayet stars

*Q: how could we understand this?*

## Supernova 1987A

Supernova Discovery Strategy III: get lucky!  
very nearby event goes off in modern age

**explosion:** Feb 23, 1987, in Large Magellanic Cloud (LMC)

$d_{\text{LMC}} \sim 50 \text{ kpc}$  – nearest (known) event in centuries

**spectrum:** shows **hydrogen**, thus **Type II event** → core collapse

**pre-explosion images:** progenitor  $M \sim 18 - 20 M_{\odot}$

star was blue supergiant

**explosion energy:** baryonic ejecta have  $1.4 \pm 0.6 \text{ foe}$

**compact remnant:** **no pulsar seen (yet)** → a black hole instead?

**ejecta:**  $M(\text{O}) \sim 2M_{\odot}$  observed;  $M(\text{Fe}) = 0.7M_{\odot}$

also N, Ne, Mg, Ni; also molecules and dust formation

light echoes: outburst reflections off surrounding material  
allow for 3-D reconstruction of pre-explosion environment!


# SN1987A: Light Curve

**light curve:** luminosity  $L$  vs  $t$

www: 1987A bolometric (all-wavelength) light curve

- initially, powered by thermal energy, then adiabatically cool

- after  $\sim 1$  month: powered by  $^{56}\text{Ni}$  decay:



*Q: how can you test that this is the power source?*

- really: decay to excited state  $^{56}\text{Ni} \rightarrow ^{56}\text{Co}^* \rightarrow ^{56}\text{Co}^{\text{gs}} + \gamma$   
 $^{56}\text{Co}$  de-excitation  $\gamma$ s seen at 0.847 MeV and 1.238 MeV

but: seen earlier than expected for onion-skin star

*Q: what does this mean?*

# SN 1987A Neutrino Signal

SN 1987A detected in neutrinos

first extrasolar (in fact, extragalactic!)  $\nu$ s  
birth of neutrino astrophysics

Reliable detections: water Čerenkov

- Kamiokande, Japan
- IMB, Ohio, USA

observed  $\sim 19$  neutrinos (mostly  $\bar{\nu}_e$ ) in 12 sec

www: ‘‘neutrino curve’’

detected  $\sim$  few hrs before optical signal

Q: Why?

22

Q: what info—qualitative and quantitative—do the  $\nu$ s give?

## Qualitatively

neutrino detection demonstrates basic correctness of core-collapse picture

## Quantitatively

$\nu$  time spread: probes diffusion from protoneutron star

$\nu$  flux, energies:  $\langle E_\nu \rangle^{\text{obs}} \sim 15 \text{ MeV}$

$\Rightarrow$  -neutrino energy release  $\mathcal{E}_{\bar{\nu}_e} \sim \mathcal{E}_\nu/6 \sim 8 \times 10^{52} \text{ erg}$

*Q: why divide by 6?*

$\Rightarrow \mathcal{E}_\nu \sim 4 \times 10^{53} \text{ erg}$

$\Rightarrow$  observational confirmation:

by far, most  $\Delta E$  released in  $\nu$ s

$\Rightarrow$  basic core collapse picture on firm ground!

Also: signal probes  $\nu$  & particle physics

## Nearby Supernovae: May We Have Another?

Today: ready for another SN!

for event at 10 kpc, Super-K will see  $\sim 5000$  events  
gravity waves?

candidates: Betelgeuse? Eta Carinae?

But don't get too close!

- minimum safe distance:  $\sim 8$  pc

*Q: why would this ruin your whole day?*

*Q: should we alert Homeland Security today?*



## Core-Collapse Nucleosynthesis

recall: hard/impossible for simulations  
to make imploding supernova explode

but we still want to know what nucleosynthesis to expect

ideally: have one self-consistent model

- pre-supernovae evolution
- detailed explosion
- ejected material gives nuke yields

*Q: in practice, how can we proceed?*

*Q: how to calibrate the “cheat”?*

25 *Q: which results/elements most likely reliable?*

*Q: which results/elements most uncertain?*

# Supernovas Nucleosynthesis—As Best We Can

real supernovae do explode:

- most ( $\gtrsim 90\%$ ) material ejected
- compact remnant (neutron star, black hole) left behind

nucleosynthesis simulation strategy:

pick ejecta/remnant division: “**mass cut**”

*force* ejection of region outside cut

either inject energy (“thermal bomb”)

or momentum (“piston”)

or extra neutrinos (“neutrino bomb”)

calibrate: demand blast with  $E_{\text{kin}} \sim 1$  foe

26 and ejected iron-peak match SN observation

still: uncertain! → particularly in yields of heaviest elements

## Explosive Nucleosynthesis

as shock passes thru pre-SN shells

compress, heat: explosive nucleosynthesis

burning occurs if mean reaction time  $\tau_{\text{nuke}} > \tau_{\text{hydro}}$

similar processes, products as before, but also freezeout behavior

- largest effects on inner shells/heaviest elements
- little change in outer shells

resulting ejecta:

dominated by  $\alpha$ -elements  $^{12}\text{C}$ ,  $^{16}\text{O}$ , ...,  $^{44}\text{Ca}$

and iron-peak elements

## Cosmic Core-Collapse Supernovae

supernovae are rare: MW rate  $r_{\text{SN}} \sim (1 - 3)/\text{century}$   
but the universe is big:  $N_{\text{gal}} \sim 4\pi/3 d_H^3 n_* \sim 10^9$  observable  
bright ( $L_* \sim L_{\text{MW}}$ ) galaxies out to horizon

so: all-sky supernova rate inside horizon  $\Gamma_{\text{SN}} \sim 1$  event/sec!  
more careful estimate: closer to  $\Gamma_{\text{SN}} \simeq 10$  events/sec!

*Q: what makes the careful estimate higher?*

These events are all neutrino sources!

if  $\mathcal{E}_{\nu, \text{tot}} \sim 300$  foe & mean neutrino energy  $\langle \epsilon \rangle_{\nu} \sim 3T_{\nu} \sim 15$  MeV  
then *per species*  $\mathcal{N}_{\nu} \sim 2 \times 10^{57}$  neutrinos emerge  
gives all-sky neutrino flux per species

$$F_{\nu}^{\text{DSNB}} \sim \frac{\Gamma_{\text{SN}} \mathcal{N}_{\nu}}{4\pi d_H^2} \sim 3 \text{ neutrinos cm}^{-2} \text{ s}^{-1} \quad (1)$$

28

*Q: how does this compare to solar neutrinos?*

*Q: how to detect it? what if we don't? what if we do?*

# Diffuse Supernova Neutrino Background

cosmic core-collapse SNe create diffuse neutrino background  
isotropic flux in all species (flavors and antiparticles)

at energies  $E_\nu \lesssim 10$  MeV, lost:

- for regular  $\nu_e, \nu_\mu, \nu_\tau$  signal swamped by solar  $\nu$ s
- even for  $\bar{\nu}$ , backgrounds too high (radioactivity, reactors)

## Detection Strategy:

look for  $\bar{\nu}_e$  at 10–30 MeV

- SN signal dominates sources & background in this window
- detect via  $\bar{\nu}_e p \rightarrow n e^+$ : KamLAND

*Not seen* so far:

- signal within factor  $\sim 2$  of limits  $\rightarrow$  should show up soon!
- *non*-detection sets limit on  
“invisible” SN which make only  $\nu$  and BH!
- *detected* background will *measure* invisible SN rate!